

**FEED ADDITIVES AVAILABLE AT BROWN'S FOR  
POULTRY GROW-OUT**

updated 4/8/10

<b>Product</b>	<b>Feed Rate</b>	<b>Use</b>
Amprol	0.125 - 0.025%	Coccidiosis in birds up to 6 weeks
Corid 9.6% Liquid	Use 4, 8 or 16 oz to make 50 gallon of treated water	Treatment and prevention of coccidiosis.
Avatec	68 to 113 grams / Ton	8 weeks old. For prevention of Coccidiosis
BMD (BMD 50) (Bacitracin Methylene Disalicylate)	4-50 grams/Ton 200 grams / Ton	Growth promotion and mild diarrhea control. Control of transmissible enteritis
BMD Soluble	4.1 Oz pack / gallon of stock solution makes 64-128 gallons of treated water	Necrotic & ulcerative enteritis
Clinacox (Diclazuril)	.91 g/ton (1ppm) Feed at rate of 1 lb./ton	Prevention of Coccidiosis
Chlortetracycline (Pennchlor 50)	50, 200, & 400 grams/Ton	Growth promotion mild respiratory problems, and Salmonella
Aureomycin (CTC) Soluble Powder	6.4 Oz pack makes 64 gallons of treated water	Bacterial pneumonia & enteritis
Nitarstone (Histostat)	170.1 grams / ton (.75 lbs./ Ton)	Prevention of Blackhead
Oxytetracycline (Pennox 50)	10 to 500 grams / Ton	Feed efficiency and respiratory infections including air sac
Terramycin Soluble Powder	6.4 Oz pack makes 25.5 gallons of treated water	Mycoplasma & Fowl Cholera
Oxytetracycline & Neomycin base (Neo-Terramycin)	50 or 100 grams / Ton 35 or 140 grams / Ton	Aid in prevention and treatment of bacterial Scours
Penicillin (ProPen100)	2.4 to 50 grams / Ton	Improved feed efficiency

**FEED ADDITIVES AVAILABLE AT BROWN'S FOR  
POULTRY GROW-OUT**

updated 4/8/10

<b>Product</b>	<b>Feed Rate</b>	<b>Use</b>
Sulfadimethoxine & Ormetoprim (Rofenaide 40)	1/2 LB. PER TON	Drug of choice in feed for fowl cholera and salmonella infections in young flocks. It is electrostatic and will cling to equipment. Be sure to purge feeding system after use.
Sulfadimethoxine	Water Soluble Powder 1 pack in 100 Gal.	Use for severe coccidiosis, E.Coli or salmonella outbreaks. Use with care - hard on birds' kidneys.
Sulmet (Sulfamethazine Sodium - 12.5%)	16 Oz. and 1 Gallon Liquid - Dilute at 1Oz/Gal.	Use to treat for Coccidiosis, Salmonella, and Fowl Cholera Use with care-Don't overdose
Tylosin (Tylan 40)	No Turkey Clearance 4-50 grams / Ton in Chickens 800 - 1000 gms/Ton	Feed Efficiency & Chronic Respiratory Disease
PrimaLac	1-3 lbs. per Ton	Feed grade Lactobacillus for reducing stress & improving feed intake.- Pellet Stable
Biotin	Variable	If foot pads have ulcers
Copper Sulfate	1 lb per Ton	Add at 1 lb. per ton if litter eating is a problem.
Ascorbic Acid	1 lb. per Ton or Water Soluble	Undefined stress reducer. Anti-Oxidant
Apex - Poultry	1/3 to 1 lb / Ton	Combat ulcerative enteritis
Vitamins & Eelectorlytes Plus	4 & 8 Oz pack makes 128 to 256 gallons of water	Vitamins and electrolytes to counteract periods of stress
Poultry Premix (B Vitamin Pack)	1-2 lbs. per Ton	Add if feed intake reduced by extreme heat or other stress. NOTE: Do not use with electrolyte packs in water. Excessive vitamins and minerals toxic.
Galliacid	1-3 lbs. / Ton	Coated organic acid to reduce pH of intestinal contents to

**FEED ADDITIVES AVAILABLE AT BROWN'S FOR  
POULTRY GROW-OUT**

updated 4/8/10

Product	Feed Rate	Use
		prevent growth of harmful bugs
Tramisol Lavamisole Hydrochloride	Water Soluble Wormer 1 bottle treats 5000 lbs. Body Wt.	Controls gape worm and intestinal worms including round worm
Wazine Piperazine	Water Soluble Wormer 1 oz / gallon of water	Add to water to worm birds 2-3x per flock. Must special order.
Salt and or Methionine	1-2 lbs./Ton	Will reduce feather picking in flocks with serious problems.
Grit Starter Grit Grower Grit Developer Grit	Variable	Starter - feed small amounts after 3 days of age.  Grower - offer free choice to birds if litter eating is a problem
York Calcium Chips	Variable	Use as grit and calcium additive in breeders and in older birds with weak/brittle bones.
Wormall Pellets Fenbendazole /Safeguard	1 lb. per 500 lbs. of Turkey	Worming via feed.
18% Turkey Wormer 21% Turkey Wormer Fenbendazole /Safeguard	Feed for 6 days as a complete feed.	Only available in 2 ton or larger lots. Worming via feed
Bicarb	1-2 lbs. per Ton	Add to feed in severely hot weather to improve lung capacity.
Fat or Soy Oil	Up to 100 lbs./Ton of total fats or oils	Adds energy to feeds in hot weather but pellet quality suffers. Already have 80 lbs. in your Grower, Developer and Finisher feed.
Oats	Variable - 100 to 300 lbs. / Ton	Can be used to slow growth and increase finish of birds 15 weeks and over.